



OsteoPlus (Osteogenesis Differentiation Medium)

For research use only

Catalogue number: BI-1102

Product Description

OsteoPlus is a ready-to-use medium developed for the in vitro osteogenesis differentiation of mesenchymal stem cells (MSCs) especially human mesenchymal stem cells. When used as directed, this media will support differentiation of these stem cells into osteogenic or bone producing cells. The medium contains all reagents for induction of osteogenesis pathways in MSCs to generate osteocytes.

Important information

- OsteoPlus is stable up to 1 month at 2 to 8°C. For long term storage, you need to freeze the medium in small aliquots. However, freezing reduces percent of differentiation.
- Do not freeze and thaw the medium.
- The medium is enriched with Glutamax.
- The medium should be clear red in color. Do not use if the color is changes.

Passaging and preparation of MSCs

1. Observe cell monolayer from basal cultures expanded in standard growth medium (DMEM (BI-1004) supplemented with 10% FBS (BI-1201) is recommended) to ensure mid-log growth phase confluence (60–80%). Aspirate medium and floating cells from culture flask and discard. Do not let passaged MSCs become completely confluent, as it can reduce the multipotency of MSCs.
2. Add 5–10 mL DPBS (BI-1401). Gently rinse cell monolayer.
3. Remove DPBS, add an adequate amount of pre-warmed trypsin from Bioidea (BI-1603 or BI-1604) to the flask and completely coat culture surface. Incubate for 5–8 minutes at 36°C to 38°C or until cells have fully detached. Overexposure to trypsin will lead to reduced MSC viability and expansion.
4. Gently pipet detached cells into a single cell solution and verify by inverted microscope.
5. Remove cell suspension from the flask, transfer to a centrifuge tube.
6. Pellet cells at 100 × g for 5–10 minutes.
7. Determine cell viability and total cell density using Trypan Blue Stain (BI-1803).
8. Re-suspend the pellet in an appropriate volume of pre-warmed MSC Growth Medium (DMEM + 10% FBS is recommended).
9. Incubate the cells in MSC Growth Medium at 36°C to 38°C in a humidified atmosphere of 4–6% CO₂ for a minimum of 2 hours up to 4 days. Inoculate plates at passage of T25 flask at approximately 1 × 10⁴ cells/cm² for low passage cells (strongly recommended), or 2 × 10⁴ cells/cm² for high passage cells in standard growth medium (DMEM + 10% FBS is recommended). Continuously passaged MSCs will gradually lose their multipotency with increased passage number (>10 passages). Use one of the following quantities and plates:
 - 2 ml growth medium per well for 2-well culture slides, OR
 - 1 ml per well for 12-well plates, OR
 - 0.5 ml per well for 24-well plates, OR
 - 200 µl for 48-well plates.

Note

For subsequent spectrophotometric assay, leave triplicate wells empty for blanks, and triplicate wells of each ASC strain to be tested in non-induction medium.



Osteogenesis differentiation

10. Examine cells every day to ensure mid-log growth phase confluence (60–80%), or are just confluent, remove complete growth medium and replace replicate wells with pre-warmed OsteoPlus Medium (BI-1102) and with control medium and continue incubation. MSCs will continue to undergo limited expansion as they differentiate under osteogenic conditions. Refeed cultures every 3–4 days. Expanding MSCs in growth medium for 2–4 days before refeeding with OsteoPlus Medium can enhance osteogenesis.
11. After specific periods of cultivation, osteogenic cultures can be processed for Alizarin Red (BI-1801) beginning at 7– 14 days, gene expression analysis, or protein detection.
NOTE: Certain cell strains might continue to grow and become over-dense under these conditions, to the point that the cell monolayer clumps or lifts off.
NOTE: Some cultures may accumulate crystals more slowly than others. If the culture appears to be differentiating, but the putative crystals are still small after 2 weeks, the culture may be re-fed with fresh OsteoPlus Medium and incubation continued for another 4 days (21 days total in osteogenic conditions). Calcium crystals can be seen as early as 6 days incubated with differentiation medium.
12. The cells are then washed once with cold DPBS. If 2-well culture slides are used, aspirate the medium and add 2 ml cold DPBS per well, or 1 ml per well for 12-well and 24-well plates (use these volumes in all steps below as well).
13. Aspirate the wash and add Formaldehyde/ Calcium solution for a minimum of 10 minutes at room temperature, or a maximum of 24 hours at 4°C. If the Alizarin Red staining cannot be done by this time, aspirate the fixative, replace with cold PBS, and store at 4°C for up to one week until ready to stain.

References

1. J Bone Miner Res. 2008 Dec 8. miR-196a Regulates Proliferation and Osteogenic Differentiation in Mesenchymal Stem Cells Derived From Human Adipose Tissue. Kim YJ, Bae SW, Yu SS, Bae YC, Jung JS.
2. Life Sci. 2008 Dec 19;83(25-26):851-8. ICAT participates in proliferation and osteogenic differentiation of human adipose tissue-derived mesenchymal stem cell. Kim YJ, Kim JT, Bae YC, Suh KT, Jung JS.

Citations

3. Davoodian, Nahid, et al. "MicroRNA-122 overexpression promotes hepatic differentiation of human adipose tissue-derived stem cells." *Journal of cellular biochemistry* 115.9 (2014): 1582-1593.
4. Fayazi, Mehri, Mojdeh Salehnia, and Saeideh Ziaei. "Differentiation of human CD146-positive endometrial stem cells to adipogenic-, osteogenic-, neural progenitor-, and glial-like cells." *In Vitro Cellular & Developmental Biology-Animal* 51.4 (2015): 408-414.
5. Allahverdi, Amir, et al. "Differentiation of human mesenchymal stem cells into insulin producing cells by using a lentiviral vector carrying PDX1." *Cell Journal (Yakhteh)* 17.2 (2015): 231.
6. Alizadeh, Effat, et al. "Up regulation of liver-enriched transcription factors HNF4a and HNF6 and liver-specific microRNA (miR-122) by inhibition of Let-7b in mesenchymal stem cells." *Chemical biology & drug design* 85.3 (2015): 268-279.
7. Arezoumand, Khatereh Saei, et al. "The emu oil emulsified in egg lecithin and butylated hydroxytoluene enhanced the proliferation, stemness gene expression, and in vitro wound healing of adipose-derived stem cells." *In Vitro Cellular & Developmental Biology-Animal* 54.3 (2018): 205-216.
8. Jalilzadeh-Tabrizi, Sepideh, et al. "A Biomimetic Emu Oil-Blended Electrospun Nanofibrous Mat for Maintaining Stemness of Adipose Tissue-Derived Stem Cells." *Biopreservation and biobanking* 16.2 (2018): 66-76.
9. Taghikani, Mohammad, and Fatemeh Eskandari. "Differentiation of Human MSCs Into Nsulin Producing Cells by Using Lentiviral Vector Carrying PDX-1."